

Cervical cancer is the 10th most common cancer and 8th most common cause of cancer deaths among women in Singapore.



HPV vaccine given to your daughter

Cervarix protects against two of the most common type of HPV that causes 70% of cervical cancer. It is offered under the school-based vaccination programme.

Other HPV vaccines available in Singapore Gardasil

Extends additional protection against two other types of HPV which generally do not cause cervical cancer but genital warts.

Gardasil 9

A newer vaccine that extends protection against additional cancer causing HPV types and is being evaluated by Ministry of Health.

HPV vaccine is safe

but some may experience the following possible side effects, as with any other vaccines.

- pain, swelling, itching, redness at the site of injection
- mild to moderate fever, headache
- muscle ache, joint pain
- nausea, abdominal pain
- feeling lightheaded, dizzy or weak

Nonetheless, they are generally mild and should resolve on their own. However, parents should seek medical advice if severe reactions occur immediately after the vaccination, or if symptoms persist.

HPV RISKS & PREVENTION



HPV stands for human papillomavirus. Out of over 100 types of HPV, 2 types are responsible for 70% of cervical cancer.

The HPV vaccination is important for your daughter, and works best when given at a younger age.

HPV vaccination

Cervical cancer





It is recommended for 9-26 years old.

HPV is a common cause of cervical cancer.





HPV vaccine is safe. It can also be taken with other vaccines.

diagnosed annually in SG*





Offers long-term protection against HPV types that cause majority of cervical cancer



Figure provided by HPB/Singapore Cancer Registry, estimated from 2011 - 2015 data*

For more information, email Contact_YPS@hpb.gov.sg.





HER FUTURE DEPENDS ON YOUR DECISION TODAY

Protect your daughter against cervical cancer with the HPV vaccination.



PROTECT HER TODAY WHAT YOU NEED TO DO



DID YOU KNOW? The human papillomavirus (HPV)

vaccination protects your child against HPV infection that can cause cervical cancer.

•••

Recommended:

Females aged 9 to 26 years old



CHECK YOUR DAUGHTER'S ELIGIBILITY UNDER THE SCHOOL-BASED PROGRAMME Offered to: Secondary 1 and 2 female students





GET THE RIGHT DOSES, AT THE RIGHT TIME.

HPB's School Health Services will work with your daughter's school to schedule the HPV vaccination.

	TWO-DOSE SCHEDULE	THREE-DOSE SCHEDULE
First dose	Age 14 years or younger	Age 15 years or older
Second dose	In the following year (or between 5 - 13 months apart)	1 - 2.5 months after the first dose
Third dose		5 - 12 months after the first dose



OPT IN FOR VACCINATION

To submit the consent for HPV vaccination for your daughter, you can choose to:



•

Scan code

Log in to the online portal the QR **OR** via **go.gov.sg/hpb-ccp** using vour SingPass

If your child has already started on a HPV vaccination series, she should complete with the same HPV vaccine type as far as possible.



HPV VACCINATION COMPLETED.